

# BOOST X-TRA

11-8-4

## FOLIAR LIQUID PLANT FOOD WITH MICRONUTRIENTS

### GUARANTEED ANALYSIS

Total Nitrogen (N).....	11.0%
1.00% Nitrate Nitrogen	
2.35% Ammoniacal Nitrogen	
7.65% Urea Nitrogen	
Available Phosphoric Acid (P2O5).....	8.0%
Soluble Potash (K2O).....	5.0%
Copper (Cu).....	0.05%
Iron (Fe).....	0.01%
Manganese (Mn).....	0.05%
Zinc (Zn).....	0.05%
Derived from Ammonium Phosphate, Potassium Nitrate, Potassium Chloride and Urea. Micronutrients derived from Boric acid, Copper Sulfate, Iron Sulfate, Manganese Sulfate, and Zinc Sulfate	
"ALSO CONTAINS NONPLANT FOOD INGREDIENTS:"	
Humic Acid (derived from Leonardite)	
Weight per gal.: 10.11 lbs. @ 68°F    1210 g/Lt @ 20°C.	

### GENERAL INFORMATION

BOOST X-TRA is formulated for field, vegetable, row, fruit, tree, vine, ornamental and turf crops.  
BOOST X-TRA may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume).  
BOOST X-TRA is compatible with most fertilizers and pesticides, however, a compatibility test is recommended if the desired combination has not been used in the past.

MANUFACTURED BY  
CUSTOM AGRICULTURAL FORMULATORS  
P.O. BOX 26104, FRESNO, CALIFORNIA 93729 800-355-4457



## DANGER

### PRECAUTIONARY STATEMENTS

**WARNING** Hazardous to humans and animals. Causes eye and skin irritation. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

### FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Rinse container; empty rinsate into spray tank. Dispose of empty container according to approved local practices. Contact county agricultural commissioner for disposal information.

### DIRECTIONS FOR USE

Shake well before using.  
Always add this product to spray tank before adding a pesticide. Do not mix with dormant oils, lime-sulfur and/or highly alkaline spray materials.

### APPLICATION RATES ALL CROPS

#### Foliar Application:

Apply 2 to 4 quarts/ acre as needed on a 10 - 14 day schedule.

When rates of over 2 quarts are required, split applications of 2 quarts or less are recommended.

Avoid foliar applications of BOOST X-TRA when plants are under stress due to lack of moisture.

#### Soil Application:

For maintenance apply 1-3 quarts per acre

For mild deficiency apply 2-6 quarts per acre

#### Aerial Application:

Apply 1 quart per 5 to 10 gallons of water per acre. A concentration of 1 quart per gallon of spray may be used

### CONDITIONS OF SALE

Always follow all directions. The manufacturer cannot control weather conditions, methods of application nor the condition of the crop at time of application. Buyer assumes all risks of use, storage or handling of this product when used in a manner inconsistent with label directions. Information regarding the contents and the levels of metals in this product is available by calling (800) 355-4457..