


CROP BUILDER 8-16-8

LIQUID PLANT FOOD



KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Rinse container; empty rinsate into spray tank. Dispose of empty container according to approved local practices. Contact county agricultural commissioner for disposal information.

CONDITIONS OF SALE

Always follow all directions. The manufacturer cannot control weather conditions, methods of application nor the condition of the crop at time of application. Buyer assumes all risks of use, storage or handling of this product when used in a manner inconsistent with label directions



8-16-8

GUARANTEED ANALYSIS:

Total Nitrogen (N).....8.0%
 4.0% Ammoniacal Nitrogen
 4.0% Urea Nitrogen

Available Phosphate (P2O5).....16.0%
 Available Potash(K2O).....8.0%
 Iron(Fe).....0.3%
 (0.3% Chelated Iron)
 Manganese(Mn).....0.05%
 (0.05% Chelated Manganese)
 Zinc(Zn).....0.05%
 (0.05%Chelated Zinc)

Weight per gal: 11.2 lb. at 68°F

Derived from: Ammonium Hydroxide, Phosphoric Acid, and Potassium Phosphate, Urea, Iron Chelate, Manganese Chelate, Zinc Chelate

Chelating Agent: Ethylenediaminetetraacetic Acid

ALSO CONTAINS NON PLANT FOOD

INGREDIENTS: 3% Organic Acids derived from Leonardite and Ascophyllum Nodosum

MANUFACTURED BY:

CUSTOM AG FORMULATORS, iNC.
 P.O. BOX 26104, FRESNO, CA 93729 800-355-4457

Net Contents: 2.5 Gals. Net Wt: 28 Lbs.

Lot No.

GENERAL INFORMATION

This product is enhanced with a biostimulant solution. CROP BUILDER is designed to be used primarily for applications on row and field crops as well as tree, vegetable, berry and vine crops. CROP BUILDER is intended to be a supplement to a regular fertilizer program. Crop Builder may be applied by ground or air and applied on both soil and foliar applications. CROP BUILDER is compatible with other fertilizers and pesticides. However, a jar test is recommended prior to mixing unknown tank combinations with CROP BUILDER.

APPLICATION RATES:

ROW CROPS: Apply 1 - 2 pints/acre at early growth stage from V-3 to V-4. Applications prior to bloom, boot, and reproductive stage is recommended.

TOBACCO:

FLUE-CURED: Apply 1-1.5 pints/acre into setter barrel. Re-apply 1-1.5 pints/acre with cutworm treatment and lay-by application. Apply 1-1.5 pints/acre at first sucker treatment and reapply with same rate at third sucker treatment.

BURLEY: Apply 1-1.5 pints/acre into setter barrel. Re-apply at 1-2.0 pints/acre foliar during mid-season growth stage. A second mid-season application may also be added to the program based on crop needs.

FIELD CROPS:

Alfalfa (hay): Apply 1-2 pints/acre 5 to 14 days after each cutting.
Alfalfa (seed): Apply 1-2 pints/acre at pre-bloom and several more times at 3-week intervals.

A Biostimulant Solution