



ZINC-RUSH

9% Zinc Fertilizer

CARBON AND AMINO BASED
ZINC NUTRIENT SOLUTION

KEEP OUT OF REACH OF CHILDREN



WARNING

PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Rinse container; empty rinsate into spray tank. Dispose of empty container according to approved local practices. Contact county agricultural commissioner for disposal information.

F1286 GUARANTEED ANALYSIS

Sulfur (S).....3.5%
Zinc (Zn).....9.0%

Derived from zinc sulfate, organic acids, and amino proteins.

Wt per gallon: 10.50 lb. @ 68°F
1.33 Kg per Ltr @ 20°C

GENERAL INFORMATION:

ZINC-RUSH is designed as a foliar spray for cotton, vegetables, trees and other crops which require supplemental zinc. It may also be used as a soil-applied product.

COMPATIBILITY:

ZINC-RUSH is compatible with most commonly used pesticides. Always check for compatibility.

MANUFACTURED BY:
CUSTOM AGRICULTURAL FORMULATORS
P.O. BOX 26104, FRESNO, CA 93729 800-355-4457

DIRECTIONS FOR USE

Apply with adequate water to provide optimal coverage.

SOIL APPLICATION: Apply 1/2 to 4 pints per acre. Rates will depend upon the level of deficiency.

FOLIAR APPLICATION: Apply alone or with other foliar sprays at a rate of 1/2 to 4 pints per acre. Rate will depend upon the level of deficiency.

SHAKE WELL BEFORE USING

CONDITIONS OF SALE

Always follow all directions. The manufacturer cannot control weather conditions, methods of application nor the condition of the crop at time of application. Buyer assumes all risks of use, storage or handling of this product when used in a manner inconsistent with label directions.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>