



Sorbi-Crop Mn 5%

A Formulation for the Correction of Mn Deficiencies in Crops

GUARANTEED ANALYSIS

Total Nitrogen (N).....	7.0%
4.6% Nitrate Nitrogen	
2.4% Urea Nitrogen	
Manganese (Mn).....	5.0%
Derived from manganese nitrate and urea.	
Wt per gal: 12.1 lb. at 68°F Grams per liter: 1450 g @20°C	

GENERAL INFORMATION

Sorbi-Crop Mn 5% is formulated with Mannitol for increased uptake in foliar applications.

Sorbi-Crop Mn 5% is designed for field, vegetable, row, fruit, tree, vine, ornamental and turf crops.

Sorbi-Crop Mn 5% may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume).

Sorbi-Crop Mn 5% may be applied alone or with fertilizers or pesticides. It can be applied either to the plant foliage or to the soil.

Sorbi-Crop Mn 5% is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

CAUTION: Avoid co-application with phosphate-containing fertilizer solutions or highly alkaline fertilizer solutions.

CONDITIONS OF SALE

The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, the manufacturer does not make nor does it authorize any agent or representative to make any warranty or representation, expressed or implied, concerning this material or the use thereof except in conformity with the statements on the label. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions. If these conditions are unacceptable to the buyer, he should immediately return the unopened product to the seller.

DIRECTIONS

Add Sorbi-Crop product to tank after adding other tank mix products. Fill tank 1/3 - 1/2 full. Add pesticides, adjuvants, and other tank mix products. Add Sorbi-Crop product at recommended rate. Continue filling tank with water until full and agitate while filling until product is thoroughly mixed.

Foliar Application:

Apply 1 to 2 quarts/ acre as needed on a 10 - 14 day schedule.

Avoid foliar applications when plants are under stress due to lack of moisture.

Turf Application:

For greens and tees apply 2-4 fl oz. per 1,000 sq.ft

For sod and overseeding: apply 2 fl. oz per 1000 sq. ft

For sports fields and fairway applications: Apply 2-5 fl oz per 1000 sq.ft

Aerial Application:

Apply suggested rate of this product in 5 to 20 gallons of water per acre.

Drip Irrigation Application: Apply the suggested rate of this product in multiple applications 10 - 14 days apart throughout the growing season of the crop.

MANUFACTURED BY:
CUSTOM AGRICULTURAL FORMULATORS
P.O. BOX 26104, FRESNO, CA 93729 800-355-4457

KEEP OUT OF REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes serious eye damage skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Take off all contaminated clothing and wash clothing before reuse. Seek medical attention if skin irritation persists.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all recommended governmental procedures for disposal of pesticide spray solutions. Rinse container, empty rinsate and spray tank. Contact county agricultural commissioner for container disposal information.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



Net Contents:
Lot #:

Net Weight: